## **References for Lupin Studies**

Archer, BJ et al. Effect of fat replacement by inulin or lupin-kernel fibre on sausage patty acceptability, post-meal perceptions of satiety and food intake in men. British Journal Nutrition 2004; 91: 591-599.

Bahr et al. Lupin protein positively affects plasma LDL cholesterol and LDL:HDLcholesterol ratio in hypercholesterolemic adults after four weeks of supplementation: a randomized, controlled crossover study. Nutrition Journal 2013, 12:107

Bahr M, et al. Consuming a mixed diet enriched with lupin protein beneficiallyaffects plasma lipids in hypercholesterolemic subjects: A randomized controlled trial. ClinicalNutrition 2015; 34: 7–14

Belski R, et al. Effects of lupin-enriched foods on body composition and cardiovascular disease risk factors: a 12-month randomized controlled weight loss trial. Int J Obes (Lond), 2011; **35**(6): 810-9.

Champ M, et al. Volatile Fatty Acid production from lupin meal in the caecum of the rat: the role of cell wall polysaccharides and a-galactosides. Animal Feed Science and Technology 1991; 32, 3177-183

Dove ER et al.Lupin and soya reduce glycaemia acutely in type 2 diabetes. Br J Nutr 2011; 106(7): 1045-51.

Fechner A et al. Effects of legume kernel fibres and citrus fibre putative risk factorsfor colorectal cancer: randomized, double-blind, crossover human intervention trial. Nutr J 2013; 12:101

Fornasini M et al. Hypoglycemic effect of Lupinus mutabilis in healthy volunteers and subjects with dysglycemia. Nutr Hosp 2012;27(2): 425-33.

Fornasini SalvadorMV et al. Efficacy of a Lupinus mutabilis Sweet snack as complement to conventional type 2 diabetes mellitus treatment. Nutr Hosp 2019; 36(4): 905-911.

Gullon P et al. In vitro fermentation of lupin seeds (Lupinus albus) and broad beans (Vicia faba): dynamic modulation of the intestinal microbiota and metabolomic output. Food & Function 2015; 6(10): 3316-3322.

Hall RS, Thomas SJ, Johnson SK. Australian sweet lupin flour addition reduced the glycaemic index of a white bread breakfast without affecting palatability in healthy human volunteers. Asia Pacific Journal of Clinical Nutrition 2005; 14: 91-97.

Johnson SK et al. Sensory acceptability of white bread with added Australian sweet lupin (*Lupinus angustifolius*) kernel fibre and its glycaemic and insulinaemic responses when eaten as a breakfast. J Science Food Agr 2003; 83:1366–1372

Johnson SK, Chua V, Hall RS, Baxter AL. Lupin kernel fibre foods improve bowelfunction and beneficially modify some putative faecal risk factors for colon cancer in men. British Journal of Nutrition 2006; 95(2): 372-8.

Juskiewicz J et al. Influence of pea and lupin oligosaccharides on caecal short chainfatty acids production and nitrogen excretion patterns in rats. Pol J Food NutrSci 2006; 15/56 :77–82

Keogh J, et al. *Food intake, postprandial glucose, insulin and subjective satiety responses to three different bread-based test meals.* Appetite 2011; 57 (3): 707-10.

Kohajdová, Z., Karovičová, J. and Schmidt, Š. (2011). Lupin composition and possible use in bakery - A Review. Czech Journal of Food Sciences 2011; 29 (3): 203-211.

*Kouris-Blazos A,* Belski R. Health benefits of legumes and pulses with a focus on Australian sweet lupins. Asia Pac J Clin Nutr 2016; 25 (1): 1-17

Lee YP et al. Lupin-enriched bread increases satiety and reduces energy intake acutely. American J Clinical Nutrition 2006; 84: 975-980.

Lee YP et al. Protein, Fibre and Blood Pressure: Potential Benefit of Legumes. Clinical and Experimental Pharmacology and Physiology 2008; 35: 473–476.

Lee YP. Effects of lupin kernel flour-enriched bread on blood pressure: a controlled intervention study. Am J Clin Nutr 2009; 89: 766

Lima-Cabello E et al. Narrow-leafed lupin (Lupinus angustifolius L.) beta-conglutin proteins modulate

the insulin signaling pathway as potential type 2 diabetes treatment and inflammatory-related disease

amelioration. Molecular Nutrition & Food Research, 2017. 61(5).30.

Magni C et al. Conglutin y, a lupin seed protein, binds insulin in vitro and reduces plasma glucose levels of hyperglycemic rats. J Nutr Biochem 2004; 15:646.

Marchesi M et al. Hypolipidemic and anti-atherosclerotic effects of lupin proteins in a rabbit model. Brit J Nutr 2008; 100 (4):707-10.

Martínez-Villaluenga C et al. Influence of addition of raffinose family oligosaccharides on probiotic survival in fermented milk during refrigerated storage. International Dairy Journal 2006; 16: 768-774.

Nowicka G et al. Lupin proteins in the treatment of hypercholesterolemia. Atherosclerosis Supplements 2006; 7: 477-477.

Schopen K et al. Short-Term Effects of Lupin vs. Whey Supplementation on Glucose and Insulin Responses to a Standardized Meal in a Randomized Cross-Over Trial. Frontiers in Physiology 2017. 8.

Oliveria, L, Pinho, O, Torres, D. Lupin: Nutritonal composition and applications in the food industry. Food Chemistry 2006; 98 (4): 711-719.

Pilvi TK, et al. Lupin protein attenuates the development of hypertension and normalises the vascular function of NaCl-loaded Goto-Kakizaki rats. J Physiol Pharmacol 2006; 57 (2): 167-76.

Smith SC, et al. Lupin kernel fiber consumption modifies fecal microbiota in healthy men as determined by rRNA gene fluorescent in situ hybridization. European J Nutrition 2006; 45: 335-41.

Skalkos S, Moschonis G, Thomas C, McMillan J, **Kouris-Blazos A.** Effect of Lupin-Enriched Biscuits as Substitute Mid-Meal Snacks on Post-Prandial Interstitial Glucose Excursions in Post-Surgical Hospital Patients with Type 2 Diabetes. Nutrients 2020; 12 (5) 1239.

Tapadia M., et al. Lupin seed hydrolysate promotes G-protein-coupled receptor, intracellular Ca2+ and enhanced glycolytic metabolism-mediated insulin secretion from BRIN-BD11 pancreatic beta cells. Molecular and Cellular Endocrinology 2019; 480: 83-96.

Terruzzi, I. Insulin-mimetic action of Conglutin  $\gamma$ , a lupin protein, in mouse myoblast. Nutr Metab Cardio-vasc Dis 2001; 21:197-205.

Weiise K et al. Lupin protein compared to casein lowers the LDL cholesterol:HDLcholesterolratio of hypercholesterolemic adults. European Journal of Nutrition 2010; 49: 65–71